

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 6 :</b> <b>A61K 31/00, 31/20, C07C 57/03</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 98/24422</b> <b>(43) International Publication Date:</b> 11 June 1998 (11.06.98)
<b>(21) International Application Number:</b> PCT/US97/22192 <b>(22) International Filing Date:</b> 3 December 1997 (03.12.97) <b>(30) Priority Data:</b> 08/759,496 4 December 1996 (04.12.96) US <b>(71) Applicant:</b> WISCONSIN ALUMNI RESEARCH FOUNDATION (US/US); 614 North Walnut Street, P.O. Box 7365, Madison, WI 53707-7365 (US). <b>(72) Inventors:</b> COOK, Mark, E.; 15 Kewaunee Court, Madison, WI 53705 (US). PARK, Yeonhwa; Apartment 118, 5002 Sheboygan Avenue, Madison, WI 53705 (US). PARIZA, Michael, W.; 7102 Valhalla Trail, Madison, WI 53719 (US). <b>(74) Agent:</b> BERSON, Bennett, J.; Quarles & Brady, P.O. Box 2113, Madison, WI 53701-2113 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> METHOD FOR CONTROLLING BODY FAT AND/OR BODY WEIGHT IN ANIMALS AND PHARMACEUTICAL COMPOSITIONS FOR USE THEREIN  <b>(57) Abstract</b> <p>Methods of inhibiting lipoprotein lipase and controlling the body fat and the body weight of an animal employ an effective amount of at least one 20 carbon, conjugated, unsaturated, fatty acid. Pharmaceutical compositions for use in the method are also disclosed.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MR	Mauritania	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MW	Malawi	UA	Ukraine
BR	Brazil	IL	Israel	MX	Mexico	UG	Uganda
BY	Belarus	IS	Iceland	NE	Niger	US	United States of America
CA	Canada	IT	Italy	NL	Netherlands	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NO	Norway	VN	Viet Nam
CG	Congo	KE	Kenya	NZ	New Zealand	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	PL	Poland	ZW	Zimbabwe
CI	Cote d'Ivoire	KM	Democratic People's Republic of Korea	PT	Portugal		
CM	Cameroon	KZ	Kazakhstan	RO	Romania		
CN	China	LC	Salut Lucia	RU	Russian Federation		
CU	Cuba	LI	Liechtenstein	SD	Sudan		
CZ	Czech Republic	LK	Sri Lanka	SG	Singapore		
DE	Germany	LR	Liberia				
DK	Denmark						
EE	Estonia						

WO 98/24422

PCT/US97/22192

METHOD FOR CONTROLLING BODY  
FAT AND/OR BODY WEIGHT IN ANIMALS  
AND PHARMACEUTICAL COMPOSITIONS FOR USE THEREIN

CROSS REFERENCE TO RELATED APPLICATIONS

5 Not applicable

FEDERALLY SPONSORED RESEARCH

Not applicable

FIELD OF THE INVENTION

10 The present invention generally relates to a method  
of controlling body fat and/or body weight in an animal.  
It also relates to pharmaceutical compositions for use in  
the method.

BACKGROUND OF THE INVENTION

15 In today's health conscious society there is a great  
interest in the fat content of food. There is a special  
concern about the saturated fat content of meat because  
of its alleged relationship to blood cholesterol. As a  
result, most consumers would prefer to have meats of  
lower total and saturated fat content. As a result some  
20 meats, such as beef, are declining in popularity. There  
also is a great interest in dieting and other means of  
controlling (i.e. reducing and/or maintaining) the body  
fat and/or body weight of humans.

25 There is an obvious need for both a safe and  
effective method of controlling the body fat of animals  
and for pharmaceutical compositions for use in a method  
of controlling body fat and/or body weight in humans.

WO 98/24422

PCT/US97/22192

## BRIEF SUMMARY OF THE INVENTION

It is an object to disclose a safe and effective method of controlling the body fat and/or body weight of an animal.

5 It also is an object of the present invention to disclose new pharmaceutical compositions.

We have discovered a method of controlling the body fat and/or body weight in an animal which comprises administering to said animal a safe and effective amount  
10 of a 20 carbon, conjugated, unsaturated, fatty acid, such as 11,13-eicosadienoic acid; 12,14-eicosadienoic acid; 8,11, 13-eicosatrienoic acid; 8,12,14-eicosatrienoic acid; 5,8,11,13-eicosatetraenoic acid; and 5,8,12,14-eicosatetraenoic acid; an active derivative, such as an  
15 ester and a non-toxic salt, thereof; and a mixture thereof. Our method is effective in controlling body fat and/or body weight in both mammals and avian species.

Although not all the details of how the method of the present invention controls body fat and/or body  
20 weight are known, we have discovered that the administration of the 20 carbon, conjugated, unsaturated, fatty acids and their active derivatives inhibit adipocyte lipoprotein lipase which is known to be essential for fat accumulation.

25 We have discovered novel pharmaceutical compositions comprising a pharmaceutical carrier and an the active

WO 98/24422

PCT/US97/22192

ingredient, which comprises at least one 20 carbon, conjugated, unsaturated, fatty acid or an active derivative, such as an ester or non-toxic salt, thereof, or a mixture thereof.

5 It will be apparent to those skilled in the art that the aforementioned objects and other advantages may be achieved by the practice of the present invention.

#### BRIEF DESCRIPTION OF DRAWINGS

Not applicable

#### 10 DETAILED DESCRIPTION OF THE INVENTION

The preferred compounds for use in the present invention are the compounds, c11,t13-eicosadienoic acid and t12,c14-eicosadienoic acid. These compounds can be made by the alkaline isomerization of c11,c14-  
15 eicosadienoic acid or the alkaline isomerization or enzymatic isomerization of 9,12 octadecadienoic acid followed by the enzymatic elongation of the isomerized products.

The 20 carbon conjugated unsaturated, fatty acids  
20 having three and four double bonds (ie. the eicosatrienoic acids and the eicosatetraenoic acids) can be made either by the alkaline isomerization of c11, c14-eicosadienoic acid followed by the desaturation of the c5 and/or c8 position using desaturase enzyme or by the  
25 alkaline isomerization or the enzymatic isomerization of c9, c12 octadienoic acid followed by the enzymatic

WO 98/24427

PCT/US97/22192

elongation and desaturation of the isomerized products (ie., c9, t11-octadecadienoic acid or t10, c12 octadecadienoic acid). The desired fatty acid is then isolated from the reaction mixture by conventional means.

5 In one preferred embodiment of the method of the present invention a safe and effective amount of at least one 20 carbon, conjugated, unsaturated, fatty acid or an active derivative thereof or a mixture thereof, is added to the feed of an animal.

10 In another embodiment, at least one 20 carbon conjugated unsaturated fatty acid, or an active derivative thereof, or a mixture thereof is administered to the animal as a pharmaceutical composition which contains the 20 carbon, conjugated, unsaturated, fatty  
15 acid and a pharmaceutical carrier and optionally other ingredients. The amount of the 20 carbon, conjugated, unsaturated, fatty acid which is to be administered is not critical as long as it is enough to be effective because it is are relatively non-toxic.

20 The practice of the present invention is further illustrated by the examples which follow:

WO 98/24422

PCT/US97/22192

## Example 1

PREPARATION OF CONJUGATED EICOSADIENOIC ACID  
BY ALKALI ISOMERIZATION

Propylene glycol (100 g) containing 26g KOH was put  
5 into a 4-neck round bottom flask (500 ml). The flask was  
equipped with a mechanical stirrer, a thermometer, a  
reflux condenser, and a nitrogen inlet. (The nitrogen  
introduced was first run through two oxygen traps).

Nitrogen was bubbled through the propylene glycol.  
10 The flask was placed in an oil bath and the temperature  
raised to 180° C - 190° C and held for 10 minutes.

The flask was removed from the oil bath and up to 50  
g 11,14-eicosadienoic acid was added as the mixture was  
swirled. The flask was placed in the oil bath and  
15 maintained at 190° C for 2h.

The flask then was removed from the oil bath and  
cooled to room temperature with cold tap water. Methanol  
(200 ml) was added. The solution was transferred to a 1-  
liter separatory funnel and acidified (pH < 2) with 250  
20 ml 6N HCl. After dilution with 200 ml water, the mixture  
of conjugated eicosadienoic acid isomers, which consisted  
primarily of c11,t13-eicosadienoic acid and t12, c14-  
eicosadienoic acid, was extracted with 200 ml hexane.  
The hexane extract was first washed with 30% methanol in  
25 water (3 x 200 ml) and then washed with double distilled  
water (3 x 200 ml). Anhydrous sodium sulfate was added

WO 98/24422

PCT/US97/22192

to remove water. The hexane was removed under vacuum rotoevaporation. The conjugated eicosadienoic acid was stored under argon at -20°C and the purity determined by GC/ms analysis.

- 5            Similar results are obtained using ethylene glycol and heating at 180°C to 190°C for 2 to 3 hours.

Example 2

PREPARATION OF CONJUGATED EICOSADIENOIC ACID  
USING MICROSOMAL FRACTION

- 10           A microsomal fraction was prepared from mouse liver. A liver homogenate was prepared using 1 volume of mouse liver and 3 volumes of (w/v) 0.25 M sucrose, 1 mM EDTA, 10 mM Tris Cl (pH 7.4). The mixture was centrifuged at 12,000 g for 15 min. Supernates were centrifuged at 15   100,000 g for 1 hr and rinsed once. Pellets of the microsomal fraction were resuspended before use. All the foregoing steps were performed at 4°C.

- The microsomal fraction was assayed for enzymatic activity using an assay system containing 5 mM ATP, 0.5mM 20   CoA, 5 mM MgCl<sub>2</sub>, 0.2 mM malonly CoA, 2 mU acyl CoA synthetase, 2 mM NADPH, 5 mM glutathione, 0.1 M potassium phosphate buffer (pH 7.4), 0.8 mM fatty acid-albumin complexes (0.4mM albumin), and microsomal fraction (1-2 mg as protein). The assay mixture was incubated at 37°C 25   for 4-24 hours.



WO 98/24422

PCT/US97/22192

The conjugated eicosadienoic acid was prepared by treating conjugated octadecadienoic acid synthesized by enzymatic or alkaline isomerization with the microsomal fraction at 37° C for 20-24 hours.

- 5        The conjugated eicosadienoic acid was extracted from the enzymatic reaction mixture with chloroform:methanol (2:1) after addition of an internal standard (heptadecanoic acid).

#### Example 3

#### 10        INHIBITION OF LIPASE ACTIVITY BY CONJUGATED EICOSADIENOIC ACID

Conjugated eicosadienoic acid was prepared by the method of Example 1.

- 15        3T3-L1 preadipocytes, which were purchased from the American Type Culture Collection, were cultured and differentiated as described by Frost, S.C., and Lane, M.D. (1985) *J. Biol. Chem.*, 260, 2646-2652. Fatty acid-albumin complexes were prepared as previously described by Calder, P.C., Bond, J.A., Harvey, D.J., Gordon, S.,  
20        and Newsholme, E. A. (1990) *Biochem. J.*, 269, 807-814 with slight modifications. Heparin-releasable lipoprotein lipase (10 U heparin/ml incubation medium) was measured as described by Nilsson-Ehle, P., and Schotz, M.C. (1976) *J. Lipid Res.*, 17, 536-541. Protein  
25        was determined using the method described by Lowry, O.H., Rosebrough, N.J., Farr, A.L., and Randall, R.J. (1951) *J.*

WO 98/24422

PCT/US97/22192

*Biol. Chem.*, 193, 265-275. Cells were treated with fatty acid-albumin complexes for 48 hrs. The results which clearly indicated that the lipoprotein lipase was inhibited are shown in Table 1.

- 5 Table 1. Inhibitory effect of conjugated eicosadienoic (c20:2) acid on lipoprotein lipase activity in 3T3-L1 adipocytes.

Lipoprotein Lipase Activity (mU/ min/ mg protein)	
10 Control	10.65 ± 1.11
c20:2 (100 µM)	4.84 ± 0.80
c20:2 (200 µM)	2.66 ± 0.78

#### Example 4

- Eight pigs (20 kg. body weight) are fed a standard control diet containing 0.5% corn oil and an equal number are fed an identical diet in which 0.5% of the corn oil is replaced by 0.5% of the conjugated eicosadienoic acid mixture. Diet is provided free choice every day until the pigs are 110 kg. in weight. After the feeding period the pigs are sacrificed and the fat, protein, water and ash content of the carcasses is analyzed by proximate analysis and the fat depth estimated using ultrasound. Leanness is determined by measuring the back fat at the

WO 98/24422

PCT/US97/22192

10th rib, measuring the loin eye area and the hot carcass weight. Subjective scoring is used to determine the quality grade (i.e. marbling in the muscle). It is found that the carcasses of the pigs fed the conjugated  
5 eicosadienoic acid diet contain less fat than the pigs fed the control diet.

The method of the present invention may take other forms. For example, the 20 carbon, conjugated, unsaturated, fatty acids or their active derivatives can  
10 be administered to an animal in a pharmaceutical composition, such as tablets, capsules, solutions or emulsions, which contains a safe and effective dose of the 20 carbon, conjugated, unsaturated, fatty acids or their active derivatives.

15 The animal feeds and pharmaceutical compositions for use in the method of the present invention are those containing one or more of free 20 carbon, conjugated, unsaturated, fatty acids, such as 11,13-eicosadienoic acid, 12,14-eicosadienoic acid, 8,11,13-eicosatrienoic  
20 acid, 8,12,14-eicosatrienoic acid, 5,8,11,13-eicosatetraenoic acid and 5,8,12,14-eicosatetraenoic acid, their active derivatives or mixtures thereof, in combination with a conventional animal feed, human food supplement, or a pharmaceutical diluent.

25 The term "20 carbon, conjugated, unsaturated, fatty acid" as used herein is intended to include, without

WO 98/24422

PCT/US97/22192

limitation, the eicosadienoic acids, eicosatrienoic acids and the eicosatetraenoic acids, their isomers, their active derivatives, such as esters and salts, and mixtures thereof. The non-toxic salts of the free acids  
5 may be made by reacting the free acids with a non-toxic base. The esters of the free acids, such as the triglyceride esters, may be made by conventional methods.

The preferred method of synthesizing the conjugated eicosadienoic acids is that described in Example 1.  
10 However, the acids may also be prepared from 9,12-octadecadienoic acid by the action of an isomerase from a microorganism (e.g. Butyrivibrio fibrisolvens) in combination with a crude, purified or cloned elongase and a desaturase from human or other animal tissue or one  
15 expressed by bacteria, yeast or plants.

The exact amount of the active form of the 20 carbon, conjugated, unsaturated, fatty acid to be administered, of course, depends upon the animal, the active form employed, and the route of administration.  
20 However, generally it will be an amount ranging from about 0.0001 g/kg about 1 g/kg of the animals body weight.

Generally, the amount of the active form of the 20 carbon, conjugated, unsaturated, fatty acid employed as  
25 the active ingredient for a pharmaceutical for humans will range from about 100 parts per million (ppm) to

WO 98/24422

PCT/US97/22192

will range from about 100 parts per million (ppm) to about 10,000 ppm of the human's diet. However, the upper limit of the amount to be employed is not critical because these acids are relatively non-toxic. The daily dosage of the active ingredient for both reducing and maintaining body fat and body weight will normally be equal to from about 100 mg to about 20,000 mg of the free acid.

The pharmaceutical compositions of the present invention contain the active ingredient in combination with a pharmaceutical carrier. When the compositions are intended for oral administration the carrier can be one or more solid diluents, such as lactose or starch; if the composition is a capsule or liquid the carrier can be a vegetable oil. When the compositions are solutions or suspensions intended for parenteral administration the preferred carrier will be a liquid suitable for injection.

A representative pharmaceutical tablet has the following formula:

Conjugated Eicosadienoic Acid Mixture of Example 1  
(calculated as free acids) 600 mg  
Microcrystalline cellulose, sodium starch glycolate,  
corn starch, hydrogenated vegetable oil wax,  
magnesium stearate and talc added.

WO 98/24422

PCT/US97/22192

The normal daily dosage for reducing body fat would be one to thirty tablets per day.

A representative chewable pharmaceutical wafer has the following formula:

- 5        Conjugated Eicosadienoic Acid Mixture of Example 1  
         (calculated as free acids) 1000 mg  
         Added dextrose, sucrose, talc, stearic acid, mineral  
         oil, salt, and natural and artificial flavorings.  
         The normal daily dosage is one to twenty tablets a  
10      day.

A representative capsule would have the following formula:

- Conjugated Eicosadienoic Acid Mixture of Example 1  
         (calculated as free acids) 600 mg  
15        Vegetable oil q.s.a.d. 1000 mg.  
         The normal daily dosage is one to twenty capsules a  
day.

- It will be readily apparent to those skilled in the  
art that a number of modifications or changes may be made  
20        without departing from the spirit and scope of the  
present invention. Therefore, the invention is only to  
be limited by the claims.

WO 98/24422

PCT/US97/22192

## CLAIMS

1. A method of controlling the body fat and body weight of an animal comprises administering to the animal a safe and effective amount of a member selected from the group consisting of at least one 20 carbon, conjugated, unsaturated, fatty acid; an ester thereof; a salt thereof; and, a mixture thereof.

2. The method of claim 1 in which the member is selected from the group consisting of 11,13-eicosadienoic acid; 12,14-eicosadienoic acid; 8,11,13- eicosatrienoic acid; 8,12,14-eicosatrienoic acid; 5,8,11,13- eicosatetraenoic acid and 5,8,12,14- eicosatetraenoic acid.

3. The method of claim 1 in which the member is selected from the group consisting of 11,13-eicosadienoic acid, an active ester thereof, and an active salt thereof.

4. The method of claim 1 in which the member is selected from the group consisting of 12,14-eicosadienoic acid, an active salt thereof, and an active ester thereof.

WO 98/24422

PCT/US97/22192

5. A pharmaceutical composition comprising as the active ingredient a member selected from the group consisting of a 20 carbon, conjugated, unsaturated, fatty acid; an ester thereof; and, a salt thereof; in  
5 combination with a pharmaceutical carrier.

6. The compound c11,t13-eicosadienoic acid.

7. The compound t12,c14-eicosadienoic acid.

8. A method of preparing a member selected from 11,13-eicosadienoic acid and 12,14-eicosadienoic acid, respectively, which comprises treating a member selected from 9,11-octadecadienoic acid and 10,12-octadecadienoic  
5 acid, respectively, with an enzyme until the desired eicosadienoic acid is obtained.

9. A method of preparing a member selected from 11,13-eicosadienoic acid and 12,14-eicosadienoic acid which comprises isomerizing 11,14-eicosadienoic acid under alkaline conditions.



WO 98/24422

PCT/US97/22192

10. A method of preparing a member selected from  
8,11,13-eicosatrienoic acid; 8,12,14-eicosatrinoic acid;  
5,8,11,13-eicosatetraenoic acid and 5,8,12,14-  
eicosatetraenoic acid which comprises isomerizing 11,14-  
5 eicosadienoic acid under alkaline conditions, followed by  
treatment with a desaturase enzyme and the isolation of  
the desired member from the reaction mixture.

11. A method of inhibiting lipoprotein lipase in  
adipocytes which comprises treating said adipocytes with  
a safe and effective amount of a member selected from the  
group consisting of a 20 carbon, conjugated, unsaturated,  
5 fatty acid; an ester thereof; a salt thereof; and, a  
mixture thereof.